

Casimir effect for tachyonic fields

Marcin Ostrowski*

Department of Theoretical Physics, University of Łódź, ul. Pomorska 149/153, 90-236 Łódź, Poland

In this paper we examine Casimir effect in the case of tachyonic field, which is connected with particles with negative four-momentum square ($m^2 < 0$). We consider here only the case of the one dimensional, scalar field. In order to describe tachyonic field, we use the absolute synchronization scheme preserving Lorentz invariance. The renormalized vacuum energy is calculated by means of the Abel-Plana formula. Finally, the Casimir energy and Casimir force as the functions of distance are obtained. In order to compare the resulting formula with the standard one, we calculate the Casimir energy and Casimir force for massive, scalar field ($m^2 > 0$).

I. INTRODUCTION

Tachyons are hypothetical particles which move faster than light. Recently, a number of papers was published, suggesting that tachyons can play a role in cosmology, as well as in neutrino physics [1, 2]. In order to describe the kinematics of this kind of particles, formalism based on the absolute synchronization scheme should be used. Now, we briefly describe main results related to the absolute synchronization scheme. Derivation of these results and exhaustive description can be found in [3, 4].

The main idea is based on the well known fact, that the definition of the time coordinate depends on the procedure used to synchronize clocks. Furthermore, the form of Lorentz transformations depends on the synchronization scheme. In absolute synchronization a new concept, the preferred frame arise. It is intimately connected with tachyons [5]. Possibly, this distinguished inertial frame can be identified with cosmic background radiation frame.

In the formalism presented here, the transformation of coordinates between inertial frames takes the following form:

$$x'(u') = D(\Lambda, u)x(u), \quad (1)$$

where Λ is an element of Lorentz group, u is the four-velocity of the preferred frame, and $D(\Lambda, u)$ is a matrix depending on Λ and u .

Transformation law for the four-velocity of the preferred frame, which follows from Eq. (1), is defined by

$$u' = D(\Lambda, u)u. \quad (2)$$

Matrix $D(\Lambda, u)$ for any rotation R is given by

$$D(R, u) = \begin{pmatrix} 1 & 0 \\ 0 & R \end{pmatrix}, \quad (3)$$

while for the boosts w it takes the form

$$D(w, u) = \begin{pmatrix} \frac{1}{w^0} & 0 \\ -\mathbf{w} & I + \frac{\mathbf{w} \otimes \mathbf{w}^T}{1 + \sqrt{1 + \mathbf{w}^2}} - u^0 \mathbf{w} \otimes \mathbf{u}^T \end{pmatrix}, \quad (4)$$

where w is the four-velocity of the system x' with respect to the system x . Hence, in the absolute synchronization scheme,

we get the Lorentz transformation law for time coordinate in the form

$$x'^0 = x^0 / W^0. \quad (5)$$

So, in the absolute synchronization scheme, time coordinate x^0 is only rescaled by some positive factor $1/W^0$. It is the most important property, which leads to preservation of the absolute causality.

The transformation law for velocities in the absolute synchronization scheme, derived from Eqs. (1), (2) and (4) is given by

$$\mathbf{v}' = W^0 \left[\mathbf{v} + \mathbf{W} \left(\frac{\mathbf{W} \mathbf{v}}{1 + \sqrt{1 + \mathbf{W}^2}} - u^0 \mathbf{u} \mathbf{v} - 1 \right) \right]. \quad (6)$$

Contrary to the standard procedure of synchronization, the transformation law (6) is well defined not only for subluminal velocities but also for superluminal. It is continuous, and does not produce 'transcendental' tachyons moving with $|\mathbf{v}| = \infty$. This property and preservation of absolute causality make absolute synchronization scheme useful for description of tachyons.

The line element $ds^2 = g_{\mu\nu}(u)dx^\mu dx^\nu$ is invariant under Lorentz transformations if the covariant metric tensor is expressed as follows:

$$g_{\alpha\beta}(u) = \begin{pmatrix} 1 & u^0 \mathbf{u}^T \\ u^0 \mathbf{u} & -I + \mathbf{u} \otimes \mathbf{u}^T (u^0)^2 \end{pmatrix}, \quad (7)$$

where $u = (u^0, \mathbf{u})$ is four-velocity of the preferred frame. Contravariant metric tensor is given by

$$g^{\alpha\beta}(u) = \begin{pmatrix} (u^0)^2 & u^0 \mathbf{u}^T \\ u^0 \mathbf{u} & -I \end{pmatrix}. \quad (8)$$

In the case of tachyons, square of the four-momentum

$$k_\alpha k^\alpha = -m^2, \quad (9)$$

leads to the dependence of the energy k_0 on momentum $(k_1, k_2, k_3) = \mathbf{k}$ in the form

$$k_0 = (u^0)^{-1} \left[-\mathbf{u} \mathbf{k} + \sqrt{(\mathbf{u} \mathbf{k})^2 + \mathbf{k}^2 - m^2} \right], \quad (10)$$

whereas contravariant energy k^0 is

$$k^0 = u^0 \sqrt{(\mathbf{u} \mathbf{k})^2 + \mathbf{k}^2 - m^2}. \quad (11)$$

*Electronic address: m.ostrowski@merlin.fic.uni.lodz.pl

Sign of k^0 is positive and Lorentz invariant not only for massive particles but also for tachyons. Values of momentum $\mathbf{k}$ fulfil the condition in the form

$$\mathbf{k}^2 > \frac{m^2}{1 + (\mathbf{u}\mathbf{k}/|\mathbf{k}|)^2}. \quad (12)$$

Values of $\mathbf{k}$ lie outside of the oblate spheroid with half-axes m, mu^0 and with the symmetry axis parallel to $\mathbf{u}$.

A more exhaustive discussion of the absolute synchronisation scheme in the language of frame bundles is given in [4]. Another applications in physics can be found in [6] (EPR correlations) and [7] (thermodynamics).

II. ONE DIMENSIONAL TACHYONIC FIELD

The quantum description of free tachyonic field was given in [3].

In this article we consider only scalar field in one space-dimension. In this case Eqs. (10) and (11) take the form

$$k_0 = (u^0)^{-1} \left[-u^1 k_1 + \frac{1}{u^0} \sqrt{k_1^2 - (u^0 m)^2} \right], \quad (13)$$

$$k^0 = \sqrt{k_1^2 - (u^0 m)^2}. \quad (14)$$

Whereas the condition in Eq. (12) reduces to

$$k_1^2 > \frac{m^2}{1 + (u^1)^2},$$

which, with the help of the identity $(1 + u^{12})(u^0)^2 = 1$, leads to the condition

$$k_1^2 > (mu^0)^2. \quad (15)$$

The Lagrangian density of the scalar field can be written in the following form:

$$\begin{aligned} \mathcal{L} &= \frac{1}{2} (g^{\alpha\beta} \partial_\alpha \phi \partial_\beta \phi + m^2 \phi^2) = \\ &= \frac{1}{2} \left[(u^0)^2 (\partial_0 \phi)^2 + u^0 u^1 (\partial_0 \phi \partial_1 \phi + \partial_1 \phi \partial_0 \phi) + \right. \\ &\quad \left. - (\partial_1 \phi)^2 + m^2 \phi^2 \right], \end{aligned} \quad (16)$$

and canonical momentum is

$$\pi = \frac{\partial \mathcal{L}}{\partial (\partial_0 \phi)} = (u^0)^2 \partial_0 \phi + u^0 u^1 \partial_1 \phi. \quad (17)$$

Hamiltonian density is given by

$$\mathcal{H} = \frac{1}{2} \left[(u^0)^2 (\partial_0 \phi)^2 + (\partial_1 \phi)^2 - m^2 \phi^2 \right]. \quad (18)$$

Scalar field in one dimension is represented by the operator:

$$\phi(x, u) = \frac{1}{\sqrt{2\pi}} \int d\mu(k) (e^{ikx} a^+(k) + e^{-ikx} a(k)), \quad (19)$$

where $d\mu(k)$ is an invariant measure defined by

$$d\mu(k) = dk_0 dk_1 \Theta(k^0) \delta(k^2 + m^2) = dk^0 dk_1 \frac{\delta(k^0 - \omega_k)}{2\omega_k}, \quad (20)$$

while the coefficient ω_k is given by

$$\omega_k = \sqrt{k_1^2 - (u^0 m)^2}. \quad (21)$$

By means of (20), the Fourier decomposition of the field $\phi(x, u)$ takes the form

$$\phi(x, u) = \frac{1}{\sqrt{2\pi}} \int_{\Gamma} \frac{dk_1}{2\omega_k} (e^{ikx} a^+(k_1, u) + e^{-ikx} a(k_1, u)), \quad (22)$$

where the integration range Γ is determined by Eq. (15).

III. CASIMIR EFFECT FOR TACHYONIC FIELD IN THE PREFERRED FRAME

Now, let us consider the one dimensional, scalar, tachyonic field in the limited region of length a (Fig. 1) with Dirichlet boundary conditions:

$$\phi(x^0, 0) = \phi(x^0, a) = 0. \quad (23)$$

In the frame moving relative to preferred frame, energy of the particle Eq. (13) is not an even function of momentum k_1 . Hence, two waves which form the standing wave (and moving in different directions) have different length and speed. This fact causes some difficulties in calculations. In order to avoid these difficulties, in this section only Casimir energy in preferred frame ($u^0 = 1$) is computed.




FIG. 1: One dimensional field.

Taking into account the boundary condition Eq. (23) we obtain the Fourier expression of the operator ϕ such that

$$\phi(x^0, x^1) = \frac{i}{\sqrt{a}} \sum_{k_n^2 > (mu^0)^2} \frac{\sin k_n x^1}{2\omega_k} [e^{-ik_0 x^0} a_n^+ - e^{ik_0 x^0} a_n], \quad (24)$$

where $k_n = \pi n/a$. The summation in Eq. (24) is over every n which fulfill condition (15). Operators a_n and a_n^+ are annihilation and creation operators which satisfy well known commutation relations:

$$[a_n, a_{n'}] = [a_n^+, a_{n'}^+] = 0, \quad (25)$$

$$[a_n, a_{n'}^+] = 2\delta_{n,n'} \omega_n. \quad (26)$$

Hamiltonian of tachyonic field (in preferred frame) can be expressed as

$$H = \frac{1}{2} \int_0^a dx [(\partial_0 \phi)^2 + (\partial_1 \phi)^2 - m^2 \phi^2]. \quad (27)$$

On inserting operator ϕ from Eq. (24) with its derivatives into Eq. (27) and integrating with respect to variable x^1 , we find

$$H = \sum_{k_n^2 > m^2} \frac{1}{16} \omega_k^{-2} (k_0^2 - m^2 + k_n^2) (a_n^+ a_n + a_n a_n^+) = \sum_{k_n^2 > m^2} \frac{1}{8} (a_n^+ a_n + a_n a_n^+), \quad (28)$$

If we calculate vacuum energy as an expectation value of H , we get

$$E = \langle 0|H|0 \rangle = \frac{1}{2} \sum_{k_n^2 > m^2} \sqrt{k_n^2 - m^2}, \quad (29)$$

where $|0\rangle$ is a vacuum state, defined by equation $a_n|0\rangle = 0$. It is well defined and stable state (See [3]).

If we compare Eq. (29) with Eq. (13) we can see that the vacuum energy is equal to the sum of energies of each mode which satisfies boundary conditions and condition (15).

After changing the limits of summation, Eq. (29) can be written as

$$E(a) = \sum_{n=n_0}^{\infty} \sqrt{\left(\frac{\pi n}{a}\right)^2 - m^2}, \quad (30)$$

where $n_0 = [ma/\pi]$. (Function $[x]$ gives the smallest integer greater than or equal to x .) The range of summation depends on the parameter a . When it increases, the more low frequencies are rejected by function $[]$.

When a goes to infinity we have

$$\sum_{n=n_0}^{\infty} \sqrt{\left(\frac{\pi n}{a}\right)^2 - m^2} \xrightarrow{a \rightarrow \infty} \frac{a}{\pi} \int_{\frac{n_0 \pi}{a}}^{\infty} dk \sqrt{k^2 - m^2} = \frac{a}{\pi} \int_m^{\infty} dk \sqrt{k^2 - m^2}. \quad (31)$$

The last stage follows as a result of obvious equation $\lim_{x \rightarrow \infty} [x]/x = 1$.

The vacuum energy (Eq. 30), also in the case of tachyonic fields, is infinite. Therefore we introduce the renormalized ‘Casimir energy’ as a difference between vacuum energy in the two configurations shown in Fig. 2. Parameter L is the total length of considered region. In this case the Casimir energy is



FIG. 2: Two configurations of boundary conditions.

defined by

$$E_c(a) = \lim_{L \rightarrow \infty} (E_1(a) + E_2(L-a) - E_3(L)). \quad (32)$$

In order to calculate energy E_1 we use Eq. (30), while energies E_2, E_3 should be calculated with the help of Eq. (31). Hence,

$$E_c(a) = \left(\sum_{n=n_0}^{\infty} \sqrt{(\pi n/a)^2 - m^2} - \frac{a}{\pi} \int_m^{\infty} dk \sqrt{k^2 - m^2} \right). \quad (33)$$

Introducing parameter $\lambda = am/\pi$ and introducing the new variable of integration $n = ak/\pi$ we get

$$E_c(a) = \frac{\pi}{a} \left(\sum_{n=n_0}^{\infty} \sqrt{n^2 - \lambda^2} - \int_{\lambda}^{\infty} dn \sqrt{n^2 - \lambda^2} \right), \quad (34)$$

where $n_0 = [am/\pi] = [\lambda]$.

To compute the expression mentioned above we use the Abel-Plana formula

$$\sum_{n=0}^{\infty} f(n) - \int_0^{\infty} f(n) dn = \frac{1}{2} f(0) + i \int_0^{\infty} dt \frac{f(it) - f(-it)}{e^{2\pi t} - 1}. \quad (35)$$

This formula is often applied in renormalization problems. Its applications and generalizations can be found in many works. For example, we mention here [9, 11].

We can write Eq. (34) in following form:

$$E_c(a) = \frac{\pi}{a^2} \left[\sum_{n=0}^{\infty} \sqrt{(n+n_0)^2 - \lambda^2} + \int_0^{\infty} \sqrt{(n+n_0)^2 - \lambda^2} dn - \int_{\lambda}^{n_0} \sqrt{n^2 - \lambda^2} dn \right]. \quad (36)$$

Furthermore, applying of the formula (35) we finally get

$$E_c(a) = -\frac{\pi}{a} \times \left[\sqrt{2} \int_0^{\infty} dt \frac{\sqrt{(n_0^2 - \lambda^2 - t^2)^2 + 4n_0^2 t^2 - n_0^2 + \lambda^2 + t^2}}{e^{2\pi t} - 1} + \frac{1}{2} \lambda^2 \ln \left(\frac{n_0 + \sqrt{n_0^2 - \lambda^2}}{\lambda} \right) + \frac{1}{2} (n_0 - 1) \sqrt{n_0^2 - \lambda^2} \right]. \quad (37)$$

The numerical computation of the expression (34) are shown in Figs 3-6.




FIG. 3: Casimir energy of tachyonic field as a function of mass m , corresponding to $a = \pi$.

The Fig. 3 shows the Casimir energy (Eq. 34) as a function of mass m . Presented curve corresponds to distance $a = \pi$ and $u^0 = 1$. We obtain here the function piecewise differentiable, the points of indifferentiability are for integer values of parameter λ . This discontinuity of derivative is related to the existence in expressions discontinuous parameter $n_0 = [\lambda]$.




FIG. 4: Casimir energy of tachyonic field as a function of the distance a , corresponding to $m = 1$.

The Fig. 4 shows Casimir energy (Eq. 34) as a function of the parameter a (Recall that a is the distance between two points where ϕ vanishes). The presented curve corresponds to $m = 1$ and $u^0 = 1$ (preferred frame).

Since the dependence on m enters to the expression (34) only through parameter λ , therefore for other values of m the Casimir energy function will have the same character, only its horizontal and vertical scale will change.




FIG. 5: Casimir force for tachyonic field as a function of the distance a , corresponding to $m = 1$.

The Fig. 5 shows Casimir force as a function of the distance a , in preferred frame ($u^0 = 1$) for $m = 1$. The Casimir force is defined by ($F = -dE_c/da$). For $F > 0$ the force is repulsive, while for $F < 0$ force is attractive.

Energy as well as the force (Figs 4, 5), shows quasis oscillatory behaviour and changes sign many times. Force is not a differentiable function of length. There are some ‘jumps’, related to the restriction for the momentum of particles (Eq. 15). The field in a finite length a , is composed of modes, which correspond to the momenta $k_n = \pi n/a$ (Eq. 24). These momenta decrease with the increasing of distance a until they cease to satisfy condition (15) and ‘fall out’ from the energy expression.

Period of the observed quasis oscillations is equal to π/m and for masses of particles greater than 1eV has microscopic magnitude. Hence, for longer distances a ($\lambda \gg 1$) quasis oscillation might be weakly observable. So, we examine slow

changeable component of the Casimir energy, after averaging its values in single periods.




FIG. 6: Casimir energy after averaging oscillations.

The Fig. 6 shows slow changeable component of Casimir energy (as a function of λ parameter). Numerical calculations show, that presented function behaves like the power function $c\lambda^\alpha$ where $\alpha = -0.5$.

To test the solutions presented on Figs 3-6, we have computed expression in Eq. (34) by three other methods, which are not presented here. There were among others: the method of zeta function and the method of generalized Abel-Plana formula. All gave the same results.

We realize that the results presented above seem strange. Nowadays, tachyons are still hypothetical objects. Maybe it is another reason which causes that tachyons does not exist. However, this question is still open.

IV. CASIMIR EFFECT FOR MASSIVE FIELD IN THE PREFERRED FRAME

In this section we calculate Casimir energy in the case of the massive field ($m^2 > 0$) using the absolute synchronization scheme. Our purpose is the comparison of the results obtained in previous paragraph for tachyons and the well known ones for massive particles [8, 9, 10]. Analogously as on last section we consider only one dimensional, scalar field.

The formula on the square of four-momentum for massive particle

$$k_\alpha k^\alpha = +m^2, \quad (38)$$

leads to the following dependence of the energy k_0 on momentum $\underline{k}$:

$$k_0 = (u^0)^{-1} \left[-\underline{u}\underline{k} + \sqrt{(\underline{u}\underline{k})^2 + \underline{k}^2 + m^2} \right]. \quad (39)$$

Lagrangian density of field can be written as

$$\begin{aligned} \mathcal{L} &= \frac{1}{2} (g^{\alpha\beta} \partial_\alpha \phi \partial_\beta \phi - m^2 \phi^2) = \\ &= \frac{1}{2} \left[(u^0)^2 (\partial_0 \phi)^2 + u^0 u^1 (\partial_0 \phi \partial_1 \phi + \partial_1 \phi \partial_0 \phi) + \right. \\ &\quad \left. - (\partial_1 \phi)^2 - m^2 \phi^2 \right]. \end{aligned} \quad (40)$$

Scalar, one dimensional and massive field, similarly to the tachyonic case (Eq. 24), can be expressed by

$$\phi(x, u) = \frac{1}{\sqrt{2\pi}} \int d\mu'(k) (e^{ikx} a^+(k) + e^{-ikx} a(k)), \quad (41)$$

where the invariant measure $d\mu'(k)$ takes now the form

$$d\mu'(k) = dk_0 dk_1 \Theta(k^0) \delta(k^2 - m^2) = dk_0 dk_1 \frac{\delta(k^0 - \omega'_k)}{2\omega'_k}, \quad (42)$$

and the coefficient ω'_k is given by

$$\omega'_k = \sqrt{k_1^2 + (u^0 m)^2}. \quad (43)$$

(Index 'prime' is added to differentiate expressions from their tachyonic equivalents.)

The operator ϕ , with the measure $d\mu'(k)$ (Eq. 42), can be written as

$$\phi(x, u) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{dk_1}{2\omega'_k} (e^{ikx} a^+(k_1, u) + e^{-ikx} a(k_1, u)). \quad (44)$$

Contrary to the case of the tachyonic field, we have a full range of integration because the relation (38) does not lead to any restrictions for momentum values.

In this section we compare the Casimir energy for massive field with results from Section III and again we consider only Casimir effect in preferred frame ($u^0 = 1$).

Hamiltonian of the field in a finite region of length a with boundary condition (23) can be expressed by

$$H = \frac{1}{2} \int_0^a dx [(\partial_0 \phi)^2 + (\partial_1 \phi)^2 + m^2 \phi^2]. \quad (45)$$

Fourier decomposition of ϕ is defined by

$$\phi(x^0, x^1) = \frac{i}{\sqrt{a}} \sum_{n=-\infty}^{\infty} \frac{\sin k_n x^1}{2\omega'_k} [e^{-ik_0 x^0} a_n^+ - e^{ik_0 x^0} a_n], \quad (46)$$

where $k_n = \pi n/a$. Operators a_n and a_n^+ fulfill the following commutation relations:

$$[a_n, a_{n'}] = [a_n^+, a_{n'}^+] = 0, \quad (47)$$

$$[a_n, a_{n'}^+] = 2\delta_{n,n'} \omega'_n. \quad (48)$$

On inserting ϕ from Eq. (46) and its derivatives into Eq. (45) we get (after integrating with the respect to x^1) the Hamiltonian in the form:

$$\begin{aligned} H &= \sum_{n=-\infty}^{\infty} \frac{1}{16} \omega'_k{}^{-2} (k_0^2 + m^2 + k_n^2) (a_n^+ a_n + a_n a_n^+) = \\ &= \sum_{n=-\infty}^{\infty} \frac{1}{8} (a_n^+ a_n + a_n a_n^+), \end{aligned} \quad (49)$$

Now, the vacuum energy is given by

$$E = \langle 0|H|0 \rangle = \frac{1}{2} \sum_{n=-\infty}^{\infty} \sqrt{k_n^2 + m^2}, \quad (50)$$

After the change of the summation limit, we finally obtain

$$E(a) = \sum_{n=1}^{\infty} \sqrt{\left(\frac{\pi n}{a}\right)^2 + m^2}. \quad (51)$$

Renormalized Casimir energy (a difference between a vacuum energy in two configurations from Fig.1) takes the form

$$E_c(a) = \frac{\pi}{a} \left(\sum_{n=0}^{\infty} \sqrt{n^2 + \lambda^2} - \int_0^{\infty} \sqrt{n^2 + \lambda^2} \right), \quad (52)$$

where $\lambda = am/\pi$.

Using the Abel-Plana formula (35) we finally receive

$$E_c(a) = -\frac{\pi}{a} \int_{\lambda}^{\infty} \frac{\sqrt{n^2 - \lambda^2}}{\exp(2\pi n) - 1} dn. \quad (53)$$

The expression in Eq. (53) is the well known formula, describing the Casimir effect for massive field ($m^2 > 0$), in standard procedure of synchronization (See [9, 10]).

Below we present Figs 7-9, which are counterparts of the Figs 3-5. The Casimir energy as a function of m , the Casimir energy as a function of a and Casimir force are shown respectively.




FIG. 7: Casimir energy for massive field as a function of mass m , corresponding to $a = \pi$.




FIG. 8: Casimir energy for massive field as a function of the distance a , corresponding to $m = 1$.

V. CASIMIR EFFECT FOR TACHYONS IN NONTRIVIAL TOPOLOGY

In this section, we again consider the one dimensional, scalar, tachyonic field in the limited region of length a (Fig. 1).




FIG. 9: Casimir force for massive field as a function of the distance a , corresponding to $m = 1$.

But now we impose boundary conditions

$$\phi(x^0, 0) = \phi(x^0, a), \quad \partial_1(x^0, 0) = \partial_1(x^0, a), \quad (54)$$

which describe the identification of the boundary points $x^1 = 0$ and $x^1 = a$. In this section we consider Casimir energy in arbitrary frame which is moving relative to preferred frame with fourvelocity u .

Taking into account the boundary condition Eq. (54) we obtain the Fourier expression of the operator ϕ such that

$$\phi(x^0, x^1) = \frac{1}{\sqrt{a}} \sum_{k_n^2 > (mu^0)^2} \frac{1}{2\omega_k} [e^{ikx} a_n^+ + e^{-ikx} a_n], \quad (55)$$

where $k_n = 2\pi n/a$. In this case Hamiltonian of tachyonic field can be expressed as

$$H = \frac{1}{2} \int_0^a dx [(u^0)^2 (\partial_0 \phi)^2 + (\partial_1 \phi)^2 - m^2 \phi^2]. \quad (56)$$

On inserting operator ϕ from Eq. (55) with its derivatives into Eq. (56) and integrating with respect to variable x^1 , we find

$$H = \sum_{k_n^2 > (mu^0)^2} \frac{1}{8} \omega_k^{-2} ((u^0 k_0)^2 - m^2 + k_n^2) (a_n^+ a_n + a_n a_n^+). \quad (57)$$

If we calculate vacuum energy as an expectation value of H , we get

$$\begin{aligned} E &= \langle 0|H|0 \rangle = \\ &= \frac{1}{2} (u^0)^{-2} \sum_{k_n^2 > (mu^0)^2} \left(-u^0 u^1 k_n + \sqrt{k_n^2 - (u^0 m)^2} \right), \end{aligned} \quad (58)$$

After changing the limits of summation, Eq. (58) can be written as

$$E(a, u) = (u^0)^{-2} \sum_{n=n_0}^{\infty} \sqrt{\left(\frac{2\pi n}{a} \right)^2 - (mu^0)^2}, \quad (59)$$

where $n_0 = [mu^0 a / 2\pi]$. (Function $[x]$ gives the smallest integer greater than or equal to x .)

The renormalized Casimir energy (a difference between vacuum energy in the two configurations shown in Fig. 2.) is given by

$$\begin{aligned} E_c(a, u) &= (u^0)^{-2} \\ &\times \left(\sum_{n=n_0}^{\infty} \sqrt{(2\pi n/a)^2 - (mu^0)^2} - \frac{a}{2\pi} \int_{mu^0}^{\infty} dk \sqrt{k^2 - (mu^0)^2} \right). \end{aligned} \quad (60)$$

Introducing parameter $\lambda(u) = am u^0 / 2\pi$ and introducing the new variable of integration $n = ak / 2\pi$ we get

$$E_c(a, u) = \frac{2\pi}{a(u^0)^2} \left(\sum_{n=n_0}^{\infty} \sqrt{n^2 - \lambda^2(u)} - \int_{\lambda}^{\infty} dn \sqrt{n^2 - \lambda^2(u)} \right), \quad (61)$$

where $n_0 = [amu^0 / 2\pi] = [\lambda]$. The expression (61) is equal to expression (34) multiplied by factor $2(u^0)^{-2}$. Hence, in the preferred frame ($u^0 = 1$), energy (61) is two times larger than energy (34), showed in Figs 3,4. Since the dependence on u^0 enters to the expression (61) only through parameter λ and factor $\pi/a(u^0)^2$, therefore for other values of u^0 (in moving frame relative to preferred frame) the Casimir energy function will have the same character, only its horizontal and vertical scale will change.

VI. COMPARISON OF RESULTS FOR TACHYONIC AND OTHER FIELDS-CONCLUSIONS

Let us compare the results for tachyonic field (Section III) with the results obtained for the massive ones (Section IV). In the case of the massive field $m^2 > 0$ we get monotonic and differentiable dependence of energy and force (as a functions of distance). Force is always attractive and smoothly goes to zero when the distance approaches infinity.

In the tachyonic case the situation is completely different. Indeed, there is no monotonicity at obtained expressions. Energy as well as force, show quasi-oscillatory behaviour and change own sign many times. Furthermore, the force is not a differentiable function of distance. There are some 'jumps', where force goes to infinity.




FIG. 10: Casimir energy for tachyonic and massive fields in preferred frame.

Let us compare the behaviour of the Casimir energy as a function of the length a . The Fig. 10 shows Casimir energy for the tachyonic and massive fields. This figure is made in SI units for mass of particle $m = 1\text{eV}$. First (upper) curve shows Casimir energy for massive field. In this case Casimir energy decay exponentially with distance a . Second (lower) curve shows Casimir energy for the tachyonic field (averaged over oscillations). As we mentioned previously, in this case energy behaves like a power function.

For full comparison, we present on the Fig. 11 Casimir energy for massless field. This figure is made in the same units as Fig. 10. In case of the one dimensional, massless field, formula for Casimir energy takes well known form: $E_c(a) = -\pi/12a$ (See [9]).

The behaviour of Casimir energy (averaged over oscillations) for tachyonic case is more similar to massless field than to massive field. In both cases (tachyonic and massless) we have slow decay of the Casimir Energy with distance a . Con-

trary, in case of massive field, Casimir energy decay fast (exponentially) with distance a .




FIG. 11: Casimir energy for massless field in preferred frame.

-
- [1] A. Chodos, A. I. Hauser, V. A. Kosticky, *Phys. Rev. B* **150**, 431, (1985).
 - [2] J. Ciburowski, J. Rembielinski, *Eur. Phys. J. C* **8**, 157-161 (1999).
 - [3] J. Rembielinski, *Tachyons and preferred frame*, *Int. J. Mod. Phys.* **12**, 1677 (1997).
 - [4] P. Caban, J. Rembielinski, *Lorentz-covariant quantum mechanics and preferred frame*, *Phys. Rev. A* **59**, 6 (1999).
 - [5] E. C. G. Sudershan, *Tachyons and the search for a preferred frame*, North Holland Publ. Co. (1978).
 - [6] J. Rembielinski and K. A. Smolinski, *Einstein-Podolsky-Rosen correlations of spin measurements in two moving inertial frames*, *Phys. Rev. A* **66**, (2002).
 - [7] J. Rembielinski, K. A. Smolinski and G. Duniec, *Thermodynamics and preferred frame*, e-print cond-mat/0102170, (2001).
 - [8] Luscher, Symanzik and Weisz, *Nucl. Phys. B* **173**, 365, (1980).
 - [9] M. Bordag, U. Mohideen, V. M. Mostepanenko, *New Developments in the Casimir Effect*, e-print quant-ph/0106045, (2001).
 - [10] G. Plunien, B. Muller, W. Greiner, *Casimir Effect*, *Physics Reports* **134**, 87-193 (1996).
 - [11] A. A. Saharian, *The generalized Abel-Plana formula. Applications to Bessel functions and Casimir effect*, The Abdus Salam International Centre for Theoretical Physics, www.ictp.trieste.it.
 - [12] A. A. Grib, S. G. Mamayev, V. M. Mostepanenko, *Vacuum Quantum Effects in Strong Fields*, Friedmann Laboratory Publishing, St. Petersburg, (1994).
 - [13] V. M. Mostepanenko, N. N. Trunov, *The Casimir Effect and its Applications*, Clarendon Press, Oxford, (1997).
 - [14] H. B. G. Casimir, *Proc. Kon. Nederl. Akad. Wet.* **51**, (1948), 793.
 - [15] H. B. G. Casimir, D. Polder, *Phys. Rev.* **73**, (1948), 360.
 - [16] M. Planck, *Verh. d. Deutsch. Phys. Ges.* **2**, (1911), 138.
 - [17] H. Rechenberg, in M. Bordag (Ed.), *The Casimir Effect 50 Years Later*, World Scientific, Singapore, (1999), 10.
 - [18] N. N. Bogoliubov, D. V. Shirkov, *Quantum Fields*, Benjamin/Cummings, London, (1982).
 - [19] P. W. Milonni, *The Quantum Vacuum: An Introduction to Quantum Electrodynamics*, Academic Press, New York, (1994).
 - [20] V. B. Berestetskii, E. M. Lifshitz, L. P. Pitaevskii, *Quantum Electrodynamics*, Pergamon, Oxford, (1982).